

ORIGINAL INVESTIGATION

Open Access



Effect of 1,5-anhydroglucitol levels on culprit plaque rupture in diabetic patients with acute coronary syndrome

Gong Su^{1*}, Ming-Xi Gao¹, Gen-Ling Shi¹, Xi-Xi Dai¹, Wei-Feng Yao¹, Tao Zhang² and Shao-Wei Zhuang^{3*}

Abstract

Background: Postprandial hyperglycemia was reported to play a key role in established risk factors of coronary artery diseases (CAD) and cardiovascular events. Serum 1,5-anhydroglucitol (1,5-AG) levels are known to be a clinical marker of short-term postprandial glucose (PPG) excursions. Low serum 1,5-AG levels have been associated with occurrence of CAD. However, the relationship between 1,5-AG levels and coronary plaque rupture has not been fully elucidated. The aim of this study was to evaluate 1,5-AG as a predictor of coronary plaque rupture in diabetic patients with acute coronary syndrome (ACS).

Methods: A total of 144 diabetic patients with ACS were included in this study. All patients underwent intravascular ultrasound examination, which revealed 49 patients with plaque rupture and 95 patients without plaque rupture in the culprit lesion. Fasting blood glucose (FBG), hemoglobin A_{1c} (HbA_{1c}) and 1,5-AG levels were measured before coronary angiography. Fasting urinary 8-iso-prostaglandin F_{2α} (8-iso-PGF_{2α}) level was measured and corrected by creatinine clearance.

Results: Patients with ruptured plaque had significantly lower serum 1,5-AG levels, longer duration of diabetes, higher HbA_{1c} and FBG levels than patients without ruptured plaque in our study population. In multivariate analysis, low 1,5-AG levels were an independent predictor of plaque rupture (odds ratio 3.421; $P = 0.005$) in diabetic patients with ACS. The area under the receiver-operating characteristic curve for 1,5-AG (0.658, $P = 0.002$) to predict plaque rupture was superior to that for HbA_{1c} (0.587, $P = 0.087$). Levels of 1,5-AG were significantly correlated with urinary 8-iso-prostaglandin F_{2α} levels ($r = -0.234$, $P = 0.005$).

Conclusions: Serum 1,5-AG may identify high risk for coronary plaque rupture in diabetic patients with ACS, which suggests PPG excursions are related to the pathogenesis of plaque rupture in diabetes.

Keywords: 1,5-Anhydroglucitol, Plaque rupture, Acute coronary syndrome, Diabetes, Intravascular ultrasound

Background

Acute coronary syndromes (ACS), including ST-elevation myocardial infarction (STEMI) and Non-ST-elevation acute coronary syndromes (NSTEMI-ACS), are a common cause of morbidity and mortality in individuals with diabetes. Autopsy data and intravascular imaging studies have showed that ACS results from spontaneous plaque rupture or erosion and subsequent thrombosis [1–3]. A meta-analysis, based on optical coherence tomography findings, showed the rate of plaque ruptures is 70.4% in

*Correspondence: su_gong@yahoo.com; zhuangs66@yahoo.com

¹ Department of Cardiovascular Medicine, Shanghai General Hospital Baoshan Branch, No. 101 Tongtai North Road, Baoshan District, Shanghai 200940, China

³ Department of Cardiovascular Medicine, The Seventh People's Hospital, Shanghai University of Traditional Chinese Medicine, No. 358 Gaoqiaodong Road, Pudong District, Shanghai 200137, China
Full list of author information is available at the end of the article



© The Author(s) 2020. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

STEMI patients, 55.6% in NSTEMI patients and 39.1% in unstable angina patients, respectively [4]. In an analysis of lesions from patients after sudden coronary death, ruptured plaque is recognized to be responsible for the most of cases of acute coronary thrombi [5]. Diabetic patients are at a high risk for cardiovascular events for having more vulnerable features in both culprit and non-culprit lesions compare to patients without diabetes [6]. Many researchers are attempting to find out what factors could affect coronary plaque rupture in diabetic patients for preventing critical outcomes.

Serum 1,5-anhydroglucitol (1,5-AG) level is a clinical marker to better reflect short-term postprandial hyperglycemia and glycemic variability (GV) than do hemoglobin A_{1c} (HbA_{1c}) level [7]. Therefore, 1,5-AG levels may be associated with cardiovascular complications in diabetes. Indeed, some studies have reported that 1,5-AG levels are bound up with cardiovascular disease [8, 9]. Several clinical studies showed that 1,5-AG levels had utility to predict cardiovascular events in study population [10–12]. However, the association between 1,5-AG levels and coronary plaque rupture in diabetic patients with ACS is unclear. Intravascular ultrasound (IVUS) can provide detailed, high-quality tomographic images to detect plaque rupture in vivo [13]. In the present study, we investigated whether any relation exists between serum 1,5-AG level and ruptured plaque in culprit lesion identified by IVUS in diabetic patients with ACS.

Methods

Patient population and study design

This is a prospective observational study. We included 220 type 2 diabetes mellitus (T2DM) patients with ACS, who were admitted to Shanghai General Hospital Baoshan Branch and Beijing Anzhen Hospital between December 2018 and July 2019. All enrolled patients were admitted and underwent coronary angiography and IVUS in the culprit vessel. Patients with any of the following were excluded from the study: (1) totally occlusive lesions, (2) restenosis after stenting, (3) previous coronary artery bypass graft surgery, (4) severe heart failure (NYHA functional class III or above), renal failure (creatinine clearance < 30 mL/min), hepatic insufficiency, or infectious disease, (5) HbA_{1c} > 8%, (6) taking sodium-glucose cotransporter 2 inhibitor, and (7) insufficient clinical data. A total of 144 patients were included for analysis after excluding 76 patients who meeting the exclusion criteria. 49 patients had coronary plaque rupture in culprit lesion diagnosed by IVUS. ACS consisted of ST-segment elevation myocardial infarction (STEMI), non-ST-segment elevation myocardial infarction (NSTEMI) and unstable angina pectoris, which were defined according to 2013 ACCF/AHA guideline for

the management of STEMI and 2014 ACC/AHA guideline for NSTEMI-ACS. T2DM was diagnosed according to the American Diabetes Association criteria or medical history and the use of insulin or glucose-lowering medication. Hypertension was defined as systolic blood pressure \geq 140 mmHg and/or diastolic blood pressure \geq 90 mmHg or treatment with oral antihypertensive drugs. Hyperlipidemia was diagnosed according to the modified National Cholesterol Education Program-Adult Treatment Panel III. The study protocol was approved beforehand by the Medical Ethics Committee of Shanghai General Hospital Baoshan Branch and the Medical Ethics Committee of Beijing Anzhen Hospital, and the procedures followed were in accordance with the institutional guidelines. The study complied with the Declaration of Helsinki, and informed consent was obtained from all patients.

IVUS imaging protocol and analysis

All patients were performed with coronary angiography by standard Judkins technique. IVUS examination was performed using an IVUS system (iLAB™ Ultrasound Imaging System, Boston Scientific, USA) and a 40 MHz intravascular catheter (OptiCross™, Boston Scientific, USA) before any intervention. The IVUS catheter was advanced into the culprit vessel more than 10 mm beyond the culprit lesion and withdrawn at a pullback speed of 0.5 mm/s automatically. In this study, a culprit lesion was defined as the lesion related to the clinical event, as identified by both coronary angiography and electrocardiogram findings. A ruptured plaque was defined as the plaque contained a cavity that communicated with the lumen with an overlying residual fibrous cap fragment. A fragmented and loosely adherent plaque without a distinct cavity and without a fibrous cap fragment was not considered as a plaque rupture [14]. IVUS quantitative analysis was performed by two independent experienced interventional cardiologists who were blinded to the patients' clinical information according to the criteria of the American College of Cardiology Clinical Expert Consensus Document on IVUS.

Laboratory measurement

We collected blood samples and urine samples from patients after overnight fasting. Samples were stored at -80°C prior to analysis. Serum levels of 1,5-AG were measured by a colorimetric method (Nippon Kayaku, Tokyo, Japan) using a Lana 1,5-AG auto liquid automatic analyzer (JCA-BM 8060, JEOL Ltd., Tokyo, Japan). Serum concentration of hemoglobin A_{1c} (HbA_{1c}) was determined by high-performance liquid chromatographic method (Tosoh HLC-723G7; Tosoh Corporation, Tokyo, Japan). The urinary 8-iso-prostaglandin F_{2 α} (8-iso-PGF_{2 α})

levels were measured by a competitive enzyme-linked immunosorbent assay (Cayman Chemical, Ann Arbor, MI, USA) and corrected by creatinine clearance. The plasma concentration of fasting blood glucose (FBG), creatinine, total cholesterol (TC), high-density lipoprotein cholesterol (HDL-c), low-density lipoprotein cholesterol (LDL-c), triglyceride (TG), the high-sensitivity C-reactive protein (hs-CRP), N-terminal pro-brain natriuretic peptide (NT-proBNP), and troponin I (TnI) were measured. The non-HDL-c level was calculated as the TC level minus the HDL-c level.

Statistical analysis

All statistical analyses were performed by using SPSS for Windows 24.0 (SPSS Inc, Chicago, IL, USA). All data were tested for normal distribution with the Kolmogorov–Smirnov test. Data are presented as mean with standard deviation (SD) for continuous distributed variables, frequencies and percentages for categorical variables, and median with 25% and 75% percentiles for abnormal distributed parameters. Differences between two groups were assessed by using the t-tests, Chi square, and Mann–Whitney rank analysis. Correlation between continuous variables was determined by Pearson correlation coefficients. Univariate and multivariate logistic regression analyses were performed in tow models to identify independent predictors for ruptured culprit plaque in study population. 1,5-AG level was included

as a continuous variable in Model 1 and as a categorized variable (categorized into tertiles) in Model 2. The predictive value of 1,5-AG and HbA_{1c} for the presence of ruptured plaque in culprit lesion was calculated by constructing receiver-operating characteristic (ROC) curves. A value of P < 0.05 was considered statistically significant.

Results

Clinical characteristics of patients

During the study period, 220 diabetic patients with ACS underwent CAG and IVUS. We excluded 16 patients with restenosis after stenting, 15 patients with insufficient IVUS data, 5 patients with severe heart failure, 8 patients with renal failure, 14 patients without 1,5-AG data, and 18 patients with other data loss. Finally, a total of 144 patients were enrolled into the present study. Among of all subjects, 49 patients had culprit plaque rupture detected by IVUS (Rupture group), 95 patients had not (Non-rupture group) (Fig. 1). Compared to patients of non-rupture group, those patients with ruptured plaque had significantly lower 1,5-AG levels (10.5 ± 5.5 vs. 14.1 ± 7.7 $\mu\text{g/mL}$, $P=0.005$), longer duration of diabetes [median (interquartile range): 4.2 (2.0, 5.2) vs. 2.5 (1.2, 4.8) years, $P=0.009$], higher FBG (8.1 ± 3.0 vs. 7.1 ± 2.0 mmol/L, $P=0.023$), HbA_{1c} (7.2 ± 0.5 vs. $7.0 \pm 0.6\%$, $P=0.025$), and non-HDL-c [4.39 (3.63, 4.99) vs. 3.83 (3.43, 4.36) mmol/L, $P=0.006$] levels. Patients with plaque rupture also had higher hs-CRP [2.46 (1.61,




Fig. 1 Flow chart of study population

5.23) vs. 1.22 (0.84, 4.01), $P=0.014$] and urinary 8-iso-PGF_{2α} (141.9±67.2 vs. 116.1±71.6 pmol/mmolCr; $P=0.038$) levels than patients without plaque rupture. No significant differences were observed between two groups in terms of age, gender, hypertension, hyperlipidemia, body mass index (BMI), blood pressure, eGFR, and medicine treatments (Table 1).

Angiographic and IVUS results

Angiographic findings and IVUS analysis were summarized in Table 2. Culprit lesions of all 144 enrolled patients were evaluated by IVUS. Plaque rupture was observed in 49 patients (34%). There were no significant differences in culprit lesion location and three-vessel

disease between plaque rupture and non-rupture groups. IVUS data showed there were not significant differences in lesion volume, length, plaque burden, external elastic membrane cross-sectional areas, lumen cross-sectional areas, plaque plus media cross-sectional areas, and remodeling index between patients with and without plaque rupture.

Relationship between 1,5-AG level and plaque rupture

We performed univariate and multivariate analysis to determine independent predictors for plaque rupture in culprit lesion. For multivariable regression analysis in model 1, variables (age, gender, current smoking, duration of diabetes, body mass index (BMI), left ventricular

Table 1 Clinical characteristics in the study population

Variables	Rupture	Non-rupture	P value
n	49	95	
Age (years)	59±9	60±10	0.474
Males	28 (57.1)	62 (65.3)	0.340
Current smoking	34 (69.4)	51 (53.7)	0.069
Hypertension	36 (73.5)	65 (68.4)	0.531
Hyperlipidemia	32 (65.3)	58 (61.1)	0.617
Duration of diabetes (years)	4.2 (2.0, 5.2)	2.5 (1.2, 4.8)	0.009
BMI (kg/m ²)	26.4±3.6	25.6±3.5	0.110
LVEF (%)	59.5±8.3	62.2±9.2	0.092
eGFR (mL/min/1.73 m ²)	84.6±28.6	88.0±30.8	0.491
SBP (mmHg)	130±13	131±16	0.612
DBP (mmHg)	76±8	77±10	0.638
TG (mmol/L)	1.93 (1.12, 2.13)	1.72 (0.94, 1.88)	0.079
HDL-C (mmol/L)	0.96 (0.82, 1.24)	1.02 (0.91, 1.23)	0.314
Non-HDL-C (mmol/L)	4.39 (3.63, 4.99)	3.83 (3.43, 4.36)	0.006
WBC (10 ⁹ /L)	7.3±1.6	7.2±1.7	0.958
hs-CRP (mg/dL)	2.46 (1.61, 5.23)	1.22 (0.84, 4.01)	0.014
NT-proBNP	390 (155, 960)	362 (105, 890)	0.413
Tnl	1.20 (0.11, 5.12)	0.91 (0.03, 2.62)	0.228
Urinary 8-iso-PGF _{2α} (pmol/mmolCr)	141.9±67.2	116.1±71.6	0.038
FBG (mmol/L)	8.1±3.0	7.1±2.0	0.023
HbA _{1c} (%)	7.2±0.5	7.0±0.6	0.025
1,5-AG (μg/mL)	10.5±5.5	14.1±7.7	0.005
Medications on admission			
Aspirin	36 (73.5)	71 (74.7)	0.869
Statins	27 (55.1)	62 (65.3)	0.234
ACEI/ARB	25 (51.0)	52 (54.7)	0.672
Oral antidiabetic drugs	31 (63.3)	64 (67.4)	0.622
Insulin	16 (32.7)	32 (33.7)	0.901

Data are given as number (percentage) for categorical variables and mean ± standard deviation or median (IQR) for continuous variables

BMI body mass index, LVEF left ventricular ejection fraction, eGFR estimated glomerular filtration rate, SBP systolic blood pressure, DBP diastolic blood pressure, TG triglycerides, HDL-C high-density lipoprotein cholesterol, non-HDL-C non-high-density lipoprotein cholesterol, WBC white blood cell, hs-CRP high-sensitivity C-reactive protein, NT-proBNP N-terminal pro-brain natriuretic peptide, Tnl troponin I, 8-iso-PGF_{2α} 8-iso-prostaglandin F_{2α}, FBG fasting blood glucose, HbA_{1c} hemoglobin A_{1c}, 1,5-AG 1,5-anhydroglucitol, ACEI/ARB angiotensin-converting-enzyme inhibitor/angiotensin II receptor blocker

Table 2 Culprit lesion characteristics assessed by angiography and intravascular ultrasound

Variables	Rupture	Non-rupture	P value
n	49	95	
Angiographic analysis			
Culprit lesion			0.694
LM	2 (2.6)	5 (5.3)	
LAD	20 (34.2)	38 (40.4)	
LCX	9 (15.8)	16 (16.0)	
RCA	18 (47.4)	36 (38.3)	
Lesion location			0.579
Ostial	2(2.6)	6 (6.4)	
Proximal	18 (34.2)	33 (35.1)	
Mild	23 (52.6)	44 (46.8)	
Distal	6 (10.5)	12 (11.7)	
3-vessel disease	21 (42.9)	33 (34.7)	0.340
IVUS analysis			
EEM CSA (mm ²)	20.9 ± 6.4	19.4 ± 5.4	0.106
Lumen CSA (mm ²)	4.3 ± 1.4	4.2 ± 1.0	0.585
P&M CSA (mm ²)	16.6 ± 5.9	15.1 ± 5.4	0.164
Plaque burden (%)	78.6 ± 8.0	76.8 ± 8.3	0.209
Length (mm)	19.0 ± 5.4	18.6 ± 6.8	0.688
Volume (mm ³)	135.7 ± 55.3	127.4 ± 58.5	0.402
Remodeling index	1.01 ± 0.15	0.97 ± 0.17	0.196

Data are given as number (percentage) for categorical variables and mean ± standard deviation

LM left main coronary artery, LAD left anterior descending coronary artery, LCX left circumflex coronary artery, RCA right coronary artery, IVUS intravascular ultrasound, EEM external elastic membrane, CSA cross-sectional areas, P&M plaque plus media

ejection fraction (LVEF), TG, non-HDL-C, hs-CRP, urinary 8-iso-PGF_{2α}, 1,5-AG, FBG and HbA_{1c}] were entered into the univariate regression analysis, and variables with $P < 0.10$ (current smoking, duration of diabetes, LVEF, TG, non-HDL-C, urinary 8-iso-PGF_{2α}, 1,5-AG, FBG and HbA_{1c}) and forced inclusion variables that were considered as important predictors of plaque rupture or associated with 1,5-AG (age, gender, BMI) were further entered into the multivariable regression model. The result showed that 1,5-AG (OR 0.916, 95% CI 0.852–0.985, $P = 0.018$) as well as other variables (current smoking and non-HDL-C) were associated with risk of culprit plaque rupture (Table 3). In model 2, age (≥ 65 years), duration of diabetes (upper tertile, ≥ 4.8 years), BMI (> 30 kg/m²), LVEF ($< 40\%$), TG (> 1.70 mmol/L), non-HDL-C (≥ 4.1 mmol/L), hs-CRP (> 3 mg/L), 1,5-AG (lower tertile, < 9.78 μ g/mL), FBG (≥ 7.0 mmol/L), HbA_{1c} ($> 7\%$) and urinary 8-iso-PGF_{2α} (upper tertile, > 140 pmol/mmolCr) were included as categorized variables. Low level of 1,5-AG (OR 3.421, 95% CI 1.446–8.092, $P = 0.005$), current smoking (OR 3.529, 95% CI 1.292–9.638, $P = 0.014$), high level of non-HDL-c (OR 4.857,

95% CI 1.871–12.61, $P = 0.001$) and HbA_{1c} (OR 2.458, 95% CI 1.081–5.586, $P = 0.032$) were significantly associated with culprit plaque rupture in diabetic patients with ACS (Table 3). We constructed a ROC curve for predicting ruptured culprit plaque by 1,5-AG and HbA_{1c} levels in patients. The area under the ROC curve for reciprocal of 1,5-AG (0.658, 95% CI 0.563–0.752, $P = 0.002$) was significantly superior to that for HbA_{1c} (0.587, 95% CI 0.490–0.684, $P = 0.087$) (Fig. 2). The optimal cut-off value for 1,5-AG to predict culprit plaque rupture was 11.8 μ g/mL (69.4% sensitivity and 59.8% specificity).

Correlation between serum 1,5-AG level and urinary 8-iso-PGF_{2α} or other ACS biomarkers

A significant negative correlation was noted between serum level of 1,5-AG and urinary 8-iso-PGF_{2α} level ($r = -0.234$, $P = 0.005$) (Fig. 3). There was no significant correlation between the level of 1,5-AG and hs-CRP ($r = -0.116$, $P = 0.165$), TnI ($r = -0.012$, $P = 0.887$), or NT-proBNP ($r = -0.011$, $P = 0.898$). The correlation between HbA_{1c} level and urinary 8-iso-PGF_{2α} level was also not significant ($r = 0.076$, $P = 0.368$).

Discussion

Most cases of sudden cardiac death and myocardial infarction arise from thrombotic coronary occlusion following coronary plaque rupture. Diabetic patients had more plaque ruptures and thrombus than non-diabetic patients in ACS, which may be associated with the greater rates of cardiovascular events in diabetes [15]. However, available screening and diagnostic methods are insufficient to identify the victims before the event occurs. The search for noninvasive approach to detect the plaque rupture was encouraged to perform. In our present study, the principal result shows that serum 1,5-AG, as a biomarker of short-term postprandial hyperglycemia and GV, might be an important surrogate marker of coronary plaque rupture in diabetic patients with ACS.

1,5-AG levels and coronary plaque rupture

1,5-AG is a naturally occurring 1-deoxy form of glucose. As glucose levels surpass the renal threshold for glucosuria (generally around 10 mmol/L), 1,5-AG is excreted in the urine leading to a rapid reduction in serum levels [16]. Therefore, poor glycemic control is associated with low, rather than high, serum 1,5-AG levels. Unlike HbA_{1c}, 1,5-AG values are not affected by mean erythrocyte age and can accurately reflect the glycemic control in hemolytic patients [17]. On the other hand, 1,5-AG levels can also reflect the status of gestational DM in pregnant women and may anticipate the development of gestational DM [18]. In the present study, we used IVUS to identify plaque rupture

Table 3 Independent predictors for ruptured culprit plaque


Model 1	Univariate		Multivariate		Model 2	Univariate		Multivariate	
	OR (95% CI)	P value	OR (95% CI)	P value		OR (95% CI)	P value	OR (95% CI)	P value
Age	0.992 (0.958, 1.028)	0.672			Age (≥ 65 years)	0.869 (0.414, 1.824)	0.710		
Female	1.305 (0.608, 2.801)	0.494			Female	1.305 (0.608, 2.801)	0.494		
Current smoking	2.278 (0.961, 5.398)	0.062	2.647 (1.018, 6.883)	0.046	Current smoking	2.278 (0.961, 5.398)	0.062	3.529 (1.292, 9.638)	0.014
Duration of diabetes	1.059 (0.991, 1.130)	0.090			Duration of diabetes (Upper tertile, ≥ 4.8 years)	1.773 (0.851, 3.695)	0.126		
Body mass index	1.102 (0.978, 1.243)	0.111			Body mass index (> 30 kg/m ²)	1.956 (0.943, 4.055)	0.071		
LVEF	0.968 (0.932, 1.006)	0.095			LVEF (< 40%)	1.718 (0.659, 4.479)	0.268		
TG	1.353 (0.961, 1.906)	0.084			TG (> 1.70 mmol/L)	1.681 (0.810, 3.490)	0.164		
Non-HDL-C	1.904 (1.233, 2.940)	0.004	1.983 (1.236, 3.183)	0.005	non-HDL-C (≥ 4.1 mmol/L)	4.016 (1.736, 9.293)	0.001	4.857 (1.871, 12.61)	0.001
hs-CRP	1.007 (0.940, 1.080)	0.837			hs-CRP (> 3 mg/L)	1.372 (0.668, 2.817)	0.389		
1,5-AG	0.914 (0.856, 0.976)	0.007	0.916 (0.852, 0.985)	0.018	1,5-AG (Lower tertile, < 9.78 µg/mL)	3.631 (1.752, 7.524)	0.001	3.421 (1.446, 8.092)	0.005
FBG	1.176 (1.019, 1.357)	0.026			FBG (≥ 7.0 mmol/L)	1.367 (0.685, 2.729)	0.375		
HbA _{1c}	1.989 (1.083, 3.653)	0.027			HbA _{1c} (> 7%)	1.996 (0.984, 4.048)	0.055	2.458 (1.081, 5.586)	0.032
Urinary 8-iso-PGF _{2α}	1.005 (1.000, 1.010)	0.042			Urinary 8-iso-PGF _{2α} (Upper tertile, > 140 pmol/mmolCr)	2.493 (1.222, 5.085)	0.012		

Model 1: Age, duration of diabetes, body mass index, LVEF, TG, non-HDL-C, hs-CRP, 1,5-AG, FBG, HbA_{1c} and urinary 8-iso-PGF_{2α} were included as continuous variables. Model 2: All variables were included as categorized variables

LVEF left ventricular ejection fraction, TG triglycerides, non-HDL-C non-high-density lipoprotein cholesterol, hs-CRP high-sensitivity C-reactive protein, 1,5-AG 1,5-anhydroglucitol, FBG fasting blood glucose, HbA_{1c} hemoglobin A_{1c}, 8-iso-PGF_{2α} 8-iso-prostaglandin F_{2α}

in culprit lesion of diabetic patients with ACS and found that the 1,5-AG levels were significantly lower in patients with ruptured plaque than in patients without ruptured plaque. Meanwhile, patients with ruptured plaque had higher HbA_{1c}, FBG, and non-HDL-c levels compared to patients with non-ruptured plaque. This result indicates that diabetic patients with ruptured culprit plaque had worse glycometabolic and lipidemic disorders. In accordance with some recent in vivo studies [19, 20], no differences were observed for age and sex between the patients with or without culprit plaque rupture. In model 1 and model 2, univariate and multivariate logistic regression analyses showed that low 1,5-AG, high non-HDL-c levels and current smoking were independent predictors of plaque rupture of culprit lesion in diabetic patients with ACS. These results indicate that poor glycemic control and dyslipidemia

may be associated with the coronary plaque rupture in diabetes. Sheng et al. reported that increased duration of DM combined with higher HbA_{1c} levels influences culprit-plaque characteristics in patients with DM who suffer AMI [21]. In the present study, patients with ruptured plaque had longer duration of DM and higher HbA_{1c} levels than patients without ruptured plaque. The level of HbA_{1c} > 7% is associated with an increased risk of culprit plaque rupture. Moreover, the ROC curve analysis showed 1,5-AG displayed more significant value in predicting plaque rupture than HbA_{1c}. Preliminary data have shown that 1,5-AG could be expected to best reflect postprandial hyperglycemia in moderately controlled patients and was more sensitive and specific than HbA_{1c} [22]. Furthermore, as PPG increments are the major contributors to GV in T2DM, 1,5-AG may be particularly suited for monitoring




postprandial hyperglycemic excursions [23]. Unlike HbA_{1c}, 1,5-AG is not affected by hypoglycemia. As a result, 1,5-AG appears to differentiate patients with extensive PPG excursions despite having similar HbA_{1c} levels. Selvin et al. reported that patients with low 1,5-AG levels had an increased risk of coronary artery disease, stroke, heart failure, and death compared to patients with high 1,5-AG levels [10]. Takahashi et al. reported that low and exacerbated levels of 1,5-AG are associated with cardiac mortality in ACS patients [11]. The study of Fujiwara et al. showed that 1,5-AG was associated with the presence of de novo coronary artery disease in both well-controlled diabetic and non-diabetic patients [8]. Low 1,5-AG level had been found to be associated with coronary artery calcification, which may be related to the coronary plaque vulnerability [24, 25]. The current study is the first to report that 1,5-AG levels are significantly associated with coronary plaque rupture in diabetic patients with ACS. These findings may partly explain the results of previous studies that 1,5-AG levels were associated with cardiovascular outcomes and support the hypothesis that PPG excursions

is strongly associated with the atherosclerotic vulnerable plaque process.

Mechanism for PPG excursions affecting plaque rupture

Although the identified role of PPG excursions in the pathogenesis of plaque rupture has not been clarified, oxidative stress, inflammation and endothelial dysfunction may be involved in the process. It was demonstrated that glucose excursions increased oxidative stress than chronic hyperglycemia in T2DM. Ceriello et al. showed that targeting postprandial hyperglycemia has the potential to reduce oxidative stress [26]. We have recently reported that glycemic variability, a component of which is PPG excursions, was significantly correlated with oxidative stress measured as urinary 8-iso-PGF_{2α} in patients with ACS [27]. Urinary 8-iso-PGF_{2α} have been proved to be the most reliable marker to assess lipid peroxidation, which is a key mechanism for the development of atherosclerotic plaques in humans. The present study showed that serum 1,5-AG level, but not HbA_{1c}, was strongly correlated with urinary 8-iso-PGF_{2α} level in diabetic patients with ACS. This is in accordance with the



previous report of Kohata et al. that 1,5-AG is the strong correlate of oxidative stress in patients with T2DM [28], and it suggests that PPG excursions can be more important than mean glucose to induce oxidative stress in diabetes. It has been demonstrated that oxidative stress plays a key role in atherosclerotic plaque progression [29]. Our previous study showed that increased urinary 8-iso-PGF_{2α} levels were closely associated with greater absolute and percent necrotic core volumes of coronary lesions in diabetic patients [30]. In a pathological study by Nishibe et al., 8-iso-PGF_{2α} was found enriched in coronary plaque specimens especially from vulnerable patients, suggesting a crucial role of free radicals in the formation of vulnerable plaques [31]. Yura et al. reported that 8-iso-PGF_{2α} per se could stimulate endothelin-1 mRNA and protein expression in bovine aortic endothelial cells [32]. Endothelin-1 may cause the stimulation of vascular smooth muscle proliferation and formation of macrophage-rich atherosclerotic plaques. In the study of Esposito et al., the results suggested that acute hyperglycemia, and not sustained elevation of blood glucose levels, could exaggerate inflammation by an oxidative mechanism [33]. Teraguchi et al. reported that dynamic

glucose fluctuation was positively and significantly associated with CD14^{bright} CD16⁺ monocytes levels and might be related to coronary plaque rupture in patients with acute myocardial infarction [34]. All these findings suggest that postprandial hyperglycemic excursions may be involved in progression and destabilization of coronary plaques through the preferential increase in oxidative stress, proinflammatory cytokines, and endothelial dysfunction. Optimizing PPG excursions management may be helpful to prevent the rupture of coronary plaque in diabetic patients.

Study limitations

Several study limitations should be considered in the interpretation of the results. First, the sample size was relatively small, so that it may have influenced the results and the statistical analyses. Second, because we evaluated only limited patients who underwent IVUS and didn't meet any exclusion criteria, our results could have been affected by selection bias and cannot be generalized to all patients. Third, the assessment of plaque rupture was made by IVUS in this study. Although it has been demonstrated that IVUS can provide detailed,

high-quality tomographic images to detect plaque rupture, it might be likely that some plaque ruptures were undetected. More detailed plaque morphology could be obtained by combining high definition IVUS with optical coherence tomography (OCT), virtually increasing accuracy for plaque rupture detection. Finally, this is an observational study. The observational nature of analysis means that we cannot infer causality in the associations we have demonstrated. Future longitudinal and prospective studies are needed to address these issues.

Conclusions

Serum 1,5-AG displayed significant value in predicting culprit plaque rupture in diabetic patients with ACS. This suggests that PPG excursions are related to the pathogenesis of plaque rupture in diabetes. The manipulation of PPG excursions may provide a potential therapeutic target for preventing plaque rupture.

Abbreviations

1,5-AG: 1,5-Anhydroglucitol; PPG: Postprandial glucose; GV: Glycemic variability; ACS: Acute coronary syndrome; T2DM: Type 2 diabetes mellitus; IVUS: Intravascular ultrasound; 8-iso-PGF_{2α}: 8-Iso-prostaglandin F_{2α}; FBG: Fasting blood glucose; HbA_{1c}: Hemoglobin A_{1c}; BMI: Body mass index; non-HDL-C: Non-high-density lipoprotein cholesterol; hs-CRP: High-sensitivity C-reactive protein; NT-proBNP: N-terminal pro-brain natriuretic peptide; TnI: Troponin I; LVEF: Left ventricular ejection fraction.

Acknowledgements

We convey thanks to the professional technical assistance from the laboratory technicians at the Clinical Laboratory Center. The authors thank the volunteer patients for their participation, and the study nurses for their skills and devotion to the patient care.

Authors' contributions

GS and SWZ participated in the design of the study. GS and TZ participated in the exercise protocols. MXG, GLS and XXD performed laboratory measurement. WFY and TZ performed data collection. GS performed the statistical analysis and drafted the manuscript. SWZ participated in revising the manuscript. All authors read and approved the final manuscript.

Funding

This work was supported by a key grant from Outstanding Clinical Discipline Project of Shanghai Pudong (PWYgy 2018-05). This work was supported by a key grant from Beijing Health Special Foundation (JING 15-10). The funder was not involved in any study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

The study was conducted in accordance with the Declaration of Helsinki, and was approved by the Medical Ethics Committee of Shanghai General Hospital Baoshan Branch and the Medical Ethics Committee of Beijing Anzhen Hospital. All participants provided written informed consent.

Consent for publication

Not applicable.

Competing interests

The authors of the manuscript do not have any closely related papers or manuscripts that have been submitted or published elsewhere and declare that they do not have any competing interests.

Author details

¹ Department of Cardiovascular Medicine, Shanghai General Hospital Baoshan Branch, No. 101 Tongtai North Road, Baoshan District, Shanghai 200940, China. ² Center of Cardiology, Beijing Anzhen Hospital, Capital Medical University, Beijing 100029, China. ³ Department of Cardiovascular Medicine, The Seventh People's Hospital, Shanghai University of Traditional Chinese Medicine, No. 358 Gaoqiaodong Road, Pudong District, Shanghai 200137, China.

Received: 10 March 2020 Accepted: 26 May 2020

Published online: 30 May 2020

References

- Stefanadis C, Antoniou CK, Tsiachris D, Pietri P. Coronary atherosclerotic vulnerable plaque: current perspectives. *J Am Heart Assoc*. 2017;6(3):e005543.
- Saia F, Komukai K, Capodanno D, Sirbu V, Musumeci G, Bocuzzi G, et al. Eroded versus ruptured plaques at the culprit site of stemi: in vivo pathophysiological features and response to primary PCI. *JACC Cardiovasc Imaging*. 2015;8(5):566–75.
- Higuma T, Soeda T, Abe N, Yamada M, Yokoyama H, Shibutani S, et al. A combined optical coherence tomography and intravascular ultrasound study on plaque rupture, plaque erosion, and calcified nodule in patients with st-segment elevation myocardial infarction: incidence, morphologic characteristics, and outcomes after percutaneous coronary intervention. *JACC Cardiovasc Interv*. 2015;8(9):1166–76.
- Iannaccone M, Quadri G, Taha S, D'Ascenzo F, Montefusco A, Omede' P, et al. Prevalence and predictors of culprit plaque rupture at OCT in patients with coronary artery disease: a meta-analysis. *Eur Heart J Cardiovasc Imaging*. 2016;17(10):1128–37.
- Virmani R, Kolodgie FD, Burke AP, Farb A, Schwartz SM. Lessons from sudden coronary death: a comprehensive morphological classification scheme for atherosclerotic lesions. *Arterioscler Thromb Vasc Biol*. 2000;20(5):1262–75.
- Sugiyama T, Yamamoto E, Bryniarski K, Xing L, Fracassi F, Lee H, et al. Coronary plaque characteristics in patients with diabetes mellitus who presented with acute coronary syndromes. *J Am Heart Assoc*. 2018;7(14):e009245.
- Kim WJ, Park CY. 1,5-Anhydroglucitol in diabetes mellitus. *Endocrine*. 2013;43(1):33–40.
- Fujiwara T, Yoshida M, Yamada H, Tsukui T, Nakamura T, Sakakura K, et al. Lower 1,5-anhydroglucitol is associated with de novo coronary artery disease in patients at high cardiovascular risk. *Heart Vessels*. 2015;30(4):469–76.
- Ikeda N, Hara H, Hiroi Y. 1,5-Anhydro-D-glucitol predicts coronary artery disease prevalence and complexity. *J Cardiol*. 2014;64(4):297–301.
- Selvin E, Rawlings A, Lutsey P, Maruthur N, Pankow JS, Steffes M, et al. Association of 1,5-anhydroglucitol with cardiovascular disease and mortality. *Diabetes*. 2016;65(1):201–8.
- Takahashi S, Shimada K, Miyauchi K, Miyazaki T, Sai E, Ogita M, et al. Low and exacerbated levels of 1,5-anhydroglucitol are associated with cardiovascular events in patients after first-time elective percutaneous coronary intervention. *Cardiovasc Diabetol*. 2016;15(1):145.
- Ouchi S, Shimada K, Miyazaki T, Takahashi S, Sugita Y, Shimizu M, et al. Low 1,5-anhydroglucitol levels are associated with long-term cardiac mortality in acute coronary syndrome patients with hemoglobin A_{1c} levels less than 7.0. *Cardiovasc Diabetol*. 2017;16(1):151.
- Ma T, Zhou B, Hsiai TK, Shung KK. A review of intravascular ultrasound-based multimodal intravascular imaging: the synergistic approach to characterizing vulnerable plaques. *Ultrasound Imaging*. 2016;38(5):314–31.
- Tian J, Ren X, Vergallo R, Xing L, Yu H, Jia H, et al. Distinct morphological features of ruptured culprit plaque for acute coronary events compared to those with silent rupture and thin-cap fibroatheroma: a combined optical coherence tomography and intravascular ultrasound study. *J Am Coll Cardiol*. 2014;63(21):2209–16.

15. Hong YJ, Jeong MH, Choi YH, Ko JS, Lee MG, Kang WY, et al. Plaque characteristics in culprit lesions and inflammatory status in diabetic acute coronary syndrome patients. *JACC Cardiovasc Imaging*. 2009;2(3):339–49.
16. Akanuma Y, Morita M, Fukuzawa N, Yamanouchi T, Akanuma H. Urinary excretion of 1,5-anhydro- β -glucitol accompanying glucose excretion in diabetic patients. *Diabetologia*. 1988;31(11):831–5.
17. Kaniwa N, Okumiya T, Tokuhira S, Matsumura Y, Matsui H, Koga M. Hemolysis causes a decrease in HbA1c level but not in glycated albumin or 1,5-anhydroglucitol level. *Scand J Clin Lab Invest*. 2019;79(6):377–80.
18. Lorenzo-Almorós A, Hang T, Peiró C, Soriano-Guillén L, Egidio J, Tuñón J, et al. Predictive and diagnostic biomarkers for gestational diabetes and its associated metabolic and cardiovascular diseases. *Cardiovasc Diabetol*. 2019;18(1):140.
19. Kim HO, Kim CJ, Kim W, Cho JM, Soeda T, Takano M, et al. Relative risk of plaque erosion among different age and sex groups in patients with acute coronary syndrome. *J Thromb Thrombolysis*. 2020;49(3):352–9.
20. Niccoli G, Montone RA, Di Vito L, Gramagna M, Refaat H, Scalone G, et al. Plaque rupture and intact fibrous cap assessed by optical coherence tomography portend different outcomes in patients with acute coronary syndrome. *Eur Heart J*. 2015;36(22):1377–84.
21. Sheng Z, Zhou P, Liu C, Li J, Chen R, Zhou J, et al. Relationships of coronary culprit-plaque characteristics with duration of diabetes mellitus in acute myocardial infarction: an intravascular optical coherence tomography study. *Cardiovasc Diabetol*. 2019;18(1):136.
22. Dungan KM, Buse JB, Largay J, Kelly MM, Button EA, Kato S, et al. 1,5-anhydroglucitol and postprandial hyperglycemia as measured by continuous glucose monitoring system in moderately controlled patients with diabetes. *Diabetes Care*. 2006;29(6):1214–9.
23. Wang Y, Zhang YL, Wang YP, Lei CH, Sun ZL. A study on the association of serum 1,5-anhydroglucitol levels and the hyperglycaemic excursions as measured by continuous glucose monitoring system among people with type 2 diabetes in China. *Diabetes Metab Res Rev*. 2012;28(4):357–62.
24. Wada H, Dohi T, Miyauchi K, Takahashi N, Endo H, Kato Y, et al. Impact of serum 1,5-anhydro- β -glucitol level on the prediction of severe coronary artery calcification: an intravascular ultrasound study. *Cardiovasc Diabetol*. 2019;18(1):69.
25. Reith S, Milzi A, Lemma ED, Dettori R, Burgmaier K, Marx N, et al. Intrinsic calcification angle: a novel feature of the vulnerable coronary plaque in patients with type 2 diabetes: an optical coherence tomography study. *Cardiovasc Diabetol*. 2019;18(1):122.
26. Ceriello A, Monnier L, Owens D. Glycaemic variability in diabetes: clinical and therapeutic implications. *Lancet Diabetes Endocrinol*. 2019;7(3):221–30.
27. Zhang T, Su G, Mi SH, Yang HX, Xin W, Dai WL, et al. Association between blood glucose variability and the characteristics of vulnerable plaque in elderly non-ST segment elevation acute coronary syndrome patients. *Int Heart J*. 2019;60(3):569–76.
28. Kohata Y, Ohara M, Nagaïke H, Fujikawa T, Osaka N, Goto S, et al. Association of hemoglobin A_{1c}, 1,5-anhydro- β -glucitol and glycated albumin with oxidative stress in type 2 diabetes mellitus patients: a cross-sectional study. *Diabetes Ther*. 2020;11(3):655–65.
29. Pignatelli P, Menichelli D, Pastori D, Violi F. Oxidative stress and cardiovascular disease: new insights. *Kardiol Pol*. 2018;76(4):713–22.
30. Su G, Wang T, Zhang T, Yang HX, Yu SS, Dai WL, et al. Urinary 8-iso-prostaglandin F_{2a} as a risk marker for the vulnerability of culprit plaque in diabetic patients with stable coronary artery disease. *Prostaglandins Leukot Essent Fatty Acids*. 2019;140(1):11–7.
31. Nishibe A, Kijima Y, Fukunaga M, Nishiwaki N, Sakai T, Nakagawa Y, et al. Increased isoprostane content in coronary plaques obtained from vulnerable patients. *Prostaglandins Leukot Essent Fatty Acids*. 2008;78(4–5):257–63.
32. Yura T, Fukunaga M, Khan R, Nassar GN, Badr KF, Montero A. Free-radical-generated F₂-isoprostane stimulates cell proliferation and endothelin-1 expression on endothelial cells. *Kidney Int*. 1999;56(2):471–8.
33. Esposito K, Nappo F, Marfella R, Giugliano G, Giugliano F, Ciotola M, et al. Inflammatory cytokine concentrations are acutely increased by hyperglycemia in humans: role of oxidative stress. *Circulation*. 2002;106(16):2067–72.
34. Teraguchi I, Imanishi T, Ozaki Y, Tanimoto T, Orii M, Shiono Y, et al. Impact of glucose fluctuation and monocyte subsets on coronary plaque rupture. *Nutr Metab Cardiovasc Dis*. 2014;24(3):309–14.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

